

ECD), IgE (murine IgE), IgE-R (human IgE receptor  $\alpha$ -chain), MPL (human thrombopoietin receptor tyrosine kinase ECD), MusK (human muscle specific receptor tyrosine kinase ECD), NpoR (human orphan receptor NpoR ECD), Rse (human receptor tyrosine kinase, Rse, ECD), HER3 (human receptor tyrosine kinase HER3/c-erbB3 ECD), Ob-R (human leptin receptor ECD), and VEGF (human vascular endothelial growth factor) where ECD refers to the extracellular domain. The nucleotide sequence data for scFv fragments from populations of antibodies raised to each antigen was translated to derive corresponding protein sequences. The V<sub>L</sub> sequences were then compared using the program "align" with the algorithm of Feng and Doolittle (1985, 1987, 1990) to calculate the percentage identity between all pairwise combinations of chains (Feng, D.F. and Doolittle, R.F. (1985) J. Mol. Evol. 21:112-123; Feng, D.F. and Doolittle, R.F. (1987) J. Mol. Evol. 25:351-360; and Feng, D.F. and Doolittle, R.F. (1990) Methods Enzymol. 183:375-387). The percent sequence identity results of each pairwise light chain amino acid sequence comparison were arranged in matrix format (Table ~~6.1-6.15~~<sup>6</sup>).

Change(s) applied  
to document, On page 107, after line 29 and before line 30, insert Table  
6.1-6.15:  
/LAS/  
3/7/2011